

# From Archaeological Sherds to Qualitative Information for Settlement Pattern Studies

Frédérique Bertoncetto<sup>1</sup> and Laure Nuninger<sup>2</sup>

<sup>1</sup>CEPAM UMR 6130, CNRS

Valbonne, France

fberton@cepam.cnrs.fr

<sup>2</sup>Chrono-Ecologie UMR 6565, CNRS

Besançon, France

laure.nuninger@mti.univ-fcomte.fr

**Abstract.** Archaeological study of ancient settlement and territorial dynamics is usually based on field survey data, both in- and off-sites. But the spatial analysis of these surface scatters is often limited by their chronological low resolution. Using statistical and spatial analysis, the authors developed two approaches which allowed to go further in order to model settlements cycles and territorial dynamics in a long term scale with a better chronological resolution, in two areas of South-East France.

## 1. Introduction

This paper is based on data collected within the Archaeomedes project<sup>1</sup>. This European project was developed in the 90's to study the human and environmental interactions in the making of Mediterranean landscape on the long term, from Iron to Middle Ages. In this paper we will present two specific methodological experiences developed during our PhD. thesis (Bertoncetto 1999, Nuninger 2002). Our studies take place in the South-East of France, one in Languedoc close to Nîmes, and the other in Provence near Fréjus (Fig. 1). Our interest in these areas was to study settlement and territorial dynamics from the Iron Age to the late Roman period. In this paper we will focus on the earliest stage of the modelling process. It presents how we used raw data from fieldwalking to model settlement and territorial patterns.





Fig. 1. Location.

## 2. From Sherd to Spatial Analysis: Difficulties


Most of our data come from fieldwalking and can then be dated within the chronological accuracy allowed by this non-stratigraphical technique. In our studied areas, most of the sites can be dated up to the century. But it is not always possible to reach such an accuracy and a relatively large number of sites cannot be precisely dated because their material is too scarce or lack of chronological references. It is even worse for off-sites scatters since the sherds are usually shapeless and badly preserved. These dating problems may deeply alter our perception of settlement's typology and spatio-temporal distribution. Indeed, if we want to go beyond the traditional divisions in large periods (preroman / roman / late roman / medieval) in order to make a detailed analysis of the settlement's dynamics within each of them, it is necessary to use a set of data with the finest chronological resolution (i.e. one century for the roman period). This usually leads to the selection of the well-dated settlements, while the sites with the least chronological accuracy are excluded from the study. In the area of Fréjus, such a selection would have a strong quantitative effect on the database available for the study, as nearly one third of the settlements existing between the 1st and 7th century A. D. should be discarded, because they can not be precisely dated (the so-called "roman sites") (Fig. 2). It would also have a qualitative effect on the database as these chronological problems do not affect all the sites in the same way.

Whereas the big settlements which were dwelled during several centuries and usually give enough material to be accurately dated, it is not always a similar situation for the small settlements and short-lived sites, which present poorer scatters of material. At least, this also deeply alters our perception of the settlements' spatial distribution because the big sites are mainly located in the valleys (like most of the



**Fig. 2.** Two perceptions of the roman settlement pattern in the area of Fréjus, showing all the known sites dwelled between the 1st and 7th c. A. D., whatever their chronological resolution (left), and only the precisely dated sites (right).

roman villae), while the smaller sites are often situated in the highlands. Fieldwalking circumstances have an impact in this distortion since it is easier to find sherds in the plough fields of the valleys than in the mountains' woodlands. Therefore, if we only select the well-dated sites, we distort the initial set of data and create artificial spatial discontinuity, which forbid us to use spatial analysis tools to study settlements' dynamics. When it comes to study agricultural territories, we meet



**Fig. 3.** Settlements and off-site units during the Iron Age in the region of Nîmes (Vaunage, Gard): two chronological resolutions which forbid the comparison.

similar problems. The only available data are scatters of materials (off-sites), which can be interpreted as remains of ancient manuring zones (Wilkinson 1982, Hayfield 1987: 192–196, Nuninger 2002: 159–163). These archaeological data are quite precious to understand the agricultural landscape. But, while settlements can usually be dated – at least in Languedoc – century by century, these off-sites units cannot be accurately dated and they cover a long period of time. We cannot often go further from the traditional divisions in large periods, as the Iron Age for example. These different chronological resolutions make it impossible to map together settlements and off-sites units, nor to analyse their relationships, unless we consider settlements at the lowest resolution, but then we lose dynamic information (Fig. 3).

### 3. How to Produce New Information: Two Approaches


Being part of the same modelling process, which is to model territorial dynamics of ancient rural communities, the studies carried out in Languedoc and in Provence ... have developed two approaches. Each one started from different kinds of data (sites and off sites) and used different methodologies, but both produced comparable data.

#### 3.1 The Settlements' Dynamics

This approach has been developed in Provence in order to understand settlements' dynamics around the roman colony of Forum Iulii /Fréjus between the 1st and the 7th c. A. D. (Bertoncello 2003, Bertoncello in progress). All the known sites were taken into account, whatever their chronological resolution, but in order to increase the chronological accuracy of the badly dated sites, we decided to go beyond the limitations of the ceramics' typological dating techniques by paying attention to the pottery assemblages of the settlements. Assuming that contemporaneous settlements show similar assemblages of artefacts, we analysed the composition of the surface material collected on every site. All the sites were described according to the same criteria which correspond to the different types of artefacts found on the surface scatters : they include not only pottery (i.e. the different categories of amphorae, coarse and fine wares), coins, glass and metallic objects, but also the building materials and the artefacts related to craft or agricultural productions (millstones, olive or wine press counterweight, remains of pottery kiln, for example). Each of these criteria was divided into quantitative classes in order to take into account the frequency of each type of artefact on the sites. For example, the criterion "gaulish amphora" was divided into 4 categories : 1 to 4 sherds for low quantity, 5 to 10 sherds for medium quantity, 11 to 50 sherds for high quantity, more than 50 sherds for very high quantity. As for the building materials, we observed the frequency of the artefacts rather than their exact number : isolated, rare or frequent. The sites were then classified according to the composition of their material : a factor analysis followed by a cluster analysis allowed us to distinguish 15 classes of sites. Each of them presents a

specific assemblage, both regarding their qualitative (i.e. the different categories of artefact shown) and quantitative composition (the frequency of each type of artefact). The diagram of the cluster analysis clearly shows two groups of settlements (Fig. 4):

- on the left, the roman settlements, showing typical early Empire and gallo-roman assemblages of artefacts ;
- on the right, the "non-roman sites" which are, on the one hand, the pre-roman settlements (class J), and on the other hand, the late-roman sites (class O) ;
- between these two classes (J and O), the sites from classes



© f.bertoncello 2004

Fig. 4. Classification of the pottery assemblages.

K, L, M and N are not easy to date because they present very similar assemblages, mainly composed of coarse ware and few or no dating indicator.




Fig. 5. Statistical analysis of the pottery assemblages allowed to increase the chronological resolution of the badly dated roman sites and to include them in the analysis of settlement's dynamics.

The factor analysis and the classification allow us to distinguish these four classes. As shown in the diagram and factorial graph, it appears that the classes K and L are closer to the pre-roman class J, while classes M and N are closer to the late roman class O (Fig. 4). We can then assume that the settlements coming from classes K and L are more probably pre-roman, while those from classes M and N should rather be dated from Late Antiquity. This example shows how these statistical analysis can help us to increase the chronological resolution of undated sites. By this way, from the original 52 undated "roman sites" known around Fréjus, it was possible to distinguish 14 early Empire settlements and 24 late roman sites, which could then be integrated in a detailed analysis of settlements' dynamics in the area (Fig. 5).

### 3.2 Agricultural Territories

In order to finalize our perception of the ancient landscape, we can process off-sites units. With these sherds coming from domestic manure, it is possible to give a spatial pictures of agricultural fields around the settlement. The experience was carried out in Languedoc during the Iron Age. Based on non-accurate data, the aim was to create a new information with a higher chronological resolution (Nuninger 2003, Nuninger 2004). We kept all the off-sites units without any selection so as to have the best archaeological information. Then, we calculated their most likely chronological weight, i.e. their most probable existence, century by century. This processing was developed according to the frequency of each type of ceramic and considering the spatial distribution of off-site units around each settlement. Basically, for each off-site, we can define the number of sherds of each type of ceramic. Concerning the Iron Age period, only amphora can be dated, at least with a resolution of 3 or 5 centuries. Because coarse ware is used during all the period, we decided to use only amphora sherds. Around Nîmes the region was well investigated with several oppida's excavations. These investigations produced a good reference with the ratio of each type of amphora per century (Table 1.). Then, following the assumption, "spread sherds come from domestic waste", a relationship between amphora ratio discovered on the settlements and amphora collected on the fields was determined. That means that the distribution of all


Period	Etruscan amphora	Massalia amphora	Italica amphora	TOTAL
640-601	14,64 %	-	-	1,84 %
600-501	69,28 %	9,59 %	-	13,37 %
500-401	15,89 %	50,24 %	-	26,57 %
400-301	0,19 %	18,33 %	-	8,99 %
300-201	-	12,19 %	0,26 %	6,06 %
200-101	-	9,29 %	31,90 %	16,84 %
100-001	-	0,36 %	67,84 %	26,33 %
TOTAL	100 %	100 %	100 %	100 %

Table 1. Nîmes' references. Ratio per century and per type of amphora according to the frequency of sherds discovered in stratigraphical context in the region of Nîmes (Py 1990: 62).


Unit A	Etruscan amphora	Massalia amphora	Italica amphora	TOTAL
640-601	1,8	-	-	1,8
600-501	8,3	0,3	-	8,6
500-401	1,9	1,5	-	3,4
400-301	0	0,5	-	0,6
300-201	-	0,4	-	0,4
200-101	-	0,3	-	0,3
100-001	-	0	-	0
TOTAL	12	3	-	15

**Table 2.** Theoretical frequency of sherds per century according to the type of amphora : example for one unit.

types of amphora per century should be similar on the settlement locations and in the agricultural fields. Based on this relationship, the theoretical number of sherds for each off-site units was calculated according to each century. Thus, if we have three sherds of massalia amphora for unit A, during the period between 600 to 501 BC, the theoretical number of sherd is 0,3 (i.e. 3 sherds multiplied by 9,59 % which is the ratio of massalia amphora during this period according to the regional reference). Finally, in order to obtain the theoretical total of sherds per unit, the sum per century was calculated. This first operation allowed us to map the density distribution of off-site clues, century by century. These maps present a theoretical dynamic which can be compared with settlement distribution. Nevertheless, they have no value on terms of reconstitution. Their interest consist in drawing our attention to some specific areas which could have been more intensively cultivated. But their interpretation remain complex because of noise due to global approach, no selection has been done. At this step, we had to delete the units with the lowest density and to select the other off-site units with a logic of manuring area, that is to say by taking into account manuring techniques and especially the manual spreading of the manure. Ethnological knowledge stress that manure is not spread anywhere but around the settlement (Sautter 1993, Lebeau 2000). In addition, they show a strong relationship between first, the distance from the settlement to the field, then the weight of carried manure and at least the number of people who can carry it. Thus, considering this information, the settlements were classified in 3 groups and a radius was




**Fig. 6.** Selection of manuring zones.



**Fig. 7.** Spatial definition of the community land.

chosen for each of them. The radius is based on agronomical and ethnological references (Remy 1967, Barral 1968, Tissandier 1969, Mazoyer, Roudart 1997: 244). For example, a radius of 200 meters was adopted for the isolated settlement which means 12 ha of cultivated land, able to feed from 5 to 15 inhabitants (Fig. 6).

Then, the discontinuous space obtained by the spatial selection was compensated by buffering. Indeed, we assume that the exploited territory of each settlement cannot be stopped by a no man's land or empty spaces. The gaps are probably due to a lack of archaeological data. Furthermore, either the manures can be provided by other ways such as for example, green manure or animal station, or the fields can support different agricultural activities such as grazing. All the processing was systematically repeated for each settlement so as to get some homogeneous areas. For each century the resulted areas represent the theoretical community land<sup>2</sup> or permanent agricultural zones. The result cannot be considered as a landscape reconstitution, but still, it is a new way of analysing the agricultural landscape with a model based on real artefacts which is an improvement regarding to the Site Catchment Analysis model.



**Fig. 8.** The settlements' dynamics during the roman period in the area of Frejus.

#### 4. Interest and Prospects

As a result of our work on the assemblages of artefact, it was possible to have a more detailed view of the settlement's dynamics during the roman period (Fig. 8).

The 1st century AD shows the main expansion of scattered settlements in the area. They occupy all types of topographical positions : the valley, the basins and the mountains. The strong decrease of the number of sites at the end of the 2nd century AD leads to a very withdrawn settlement's pattern during the 3rd century AD. The settlements are concentrated on the biggest villas of the valley while the mountains seem to be almost deserted. The 4th and mainly the 5th century AD initiate a new dynamics with the creation of a large number of sites in the mountains. The importance of this reconquest of the mountains wouldn't have been properly recognized without the integration of unprecisely dated sites, as a lot of them appeared to be late roman. In a similar way with the analysis of off-site units, we can observe the dynamics of the exploited landscape: starting from clustered territories during 6th to 5th century BC, we can see a clear structuration of two big territories during the next centuries. At the end of the Iron Age, during the 1st century BC, it is interesting to stress some competition areas and a new small territory of conquest. The settlement patterns were modelised independently using a network organisation. But at the final step, they can be compared with the community lands (Fig. 9).




Fig. 9. The community lands' dynamics and the settlements' dynamics during the Iron Age in the area of Nîmes (Vaunage, Gard).

The comparison shows some strong relationships which can confirm part of our assumptions. Moreover, what is the most stimulating that it points out some anomalies and brings out new questions (Nuninger 2002, Nuninger 2004). Then, it is necessary to go back on the field and to work with raw data again in order to take into account these anomalies or to bring new information within the model.

The whole results give new possibilities to understand settlement cycles and territorial dynamics with a long term scale for different regions. Until now, these studies have remained experimental. But considering these preliminary results, our project is to build up an integrated model and a systematic approach on both areas.

#### Notes

- 1 Archaeomedes I (1992–1994) : “Understanding the Natural and Anthropogenic causes of soil degradation in the Mediterranean Basin” (Program Environment of the European Commission DGXII); Archaeomedes II (1996–1999) : “Policy-relevant models of the natural and anthropogenic dynamics of degradation and desertification and their spatio-temporal manifestations” (Program Environment of the European Commission DGXII).
- 2 The agricultural territory exploited by a rural community according to the definition from Leonard R. & Longbottom J. 2000, see “Terroir”: 55.

#### Acknowledgements

We wish to thank, with a specific attention, Mr Sander van der Leeuw (Archaeomedes project's coordinator) who trusted in us as young researchers and gave us a possibility to work together with a high level team.

We would like to thank the CEPAM (UMR 6130, CNRS), the laboratory of Chrono-Ecology (UMR 6565, CNRS), the MTI@SHS centre (University of Franche-Comté, France), the Regional Archaeological Service of Languedoc-Roussillon and Var (France) and the Archaeomedes Project (EU) for their funding of these works. We thank the MSH C. N. Ledoux of University of Franche-Comté (France) to support the cost of the conference for Laure Nuninger.

#### References

- Barral, H., 1968. *Tiogo (Haute-Volta). Atlas des structures agraires au sud du Sahara. 2*, Mouton/Maison des Sciences de l'Homme, Paris-La Haye.
- Bertoncello, F., 1999. Le Peuplement de la basse vallée de l'Argens et de ses marges (Var, France), de la fin de l'Age du Fer à la fin de l'Antiquité. Ph.D. thesis, defended in December 1999, University of Provence, France.
- Bertoncello, F., 2003. L'apport de la statistique descriptive à l'analyse du peuplement dans la région de Fréjus (Var) : essai de caractérisation des assemblages de mobilier. In Favory, F. and Vignot, A. (eds), *Actualité de la recherche*

- en Histoire et Archéologie agraires, Proceedings of the 5th International Conference of AGER, Besançon, France, september 19–20, 2000.* University Press of Franche-Comté, Besançon, 303–323.
- Bertoncello, F., in progress. L'occupation d'un milieu de moyenne montagne en Provence orientale : le massif des Maures (Var) du IIe s. av. n. è. au VIIe s. In Bouet, A. et Verdin, F. (eds), *Territoires et paysages. Hommages à Philippe Leveau*. Documents d'Archéologie Méridionale.
- Hayfield, C., 1987. *An Archaeological survey of the parrish of Wharram Percy. East Yorkshire: I. The evolution of the roman landscape*. Oxford. BAR, British Series; 172.
- Lebeau, R., 2000. *Les grands types de structures agraires dans le monde*. Paris, Armand Colin (U Géographie).
- Mazoyer, M., Roudart, L., 1997. *Histoire des agricultures du monde, du Néolithique à la crise contemporaine*, Paris, Seuil.
- Nuninger, L. 2002. Peuplement et territoires protohistoriques du VIIIe au Ier s. av. J.-C. en Languedoc oriental (Gard-Hérault). Ph.D. thesis, defended in December 2002, University of Franche-Comté, France:  
[http://tel.ccsd.cnrs.fr/documents/archives0/00/00/29/81/index\\_fr.html](http://tel.ccsd.cnrs.fr/documents/archives0/00/00/29/81/index_fr.html) (visited 10/07/2004).
- Nuninger, L., 2003. Approche méthodologique d'un espace agraire protohistorique: la Vaunage (Gard). *Actualité de la recherche en Histoire et Archéologie agraires, Proceedings of the 5th International Conference of AGER, Besançon, France, September 19–20, 2000.* University Press of Franche-Comté, Besançon, 365–375.
- Nuninger, L., 2004. Contribution à la modélisation des finages protohistoriques: un cas d'étude micro-régional en Languedoc oriental (Gard), *AGER* 13, 6–23.
- Py, M., 1990. *Culture, économie et société protohistoriques dans la région nîmoise*. French School of Rome, Paris-Rome, 1990.
- Remy, G., 1967. *Yobri (Haute-Volta). Atlas des structures agraires au sud du Sahara. 1*, Mouton/Maison des Sciences de l'Homme, Paris-La Haye.
- Sautter, G., 1993. *Parcours d'un géographe. Des paysages aux ethnies, de la brousse à la ville, de l'Afrique au monde*. Paris, Arguments.
- Tissandier, J., 1969. *Zengoaga (Cameroun). Atlas des structures agraires au sud du Sahara. 3*, Mouton/Maison des Sciences de l'Homme, Paris-La Haye.
- Wilkinson, T.-J., 1982. The definition of ancient manured zones by means of extensive sherd-sampling techniques. *Journal of Field Archaeology* 9–3, 323–333.